

**Station 6 - 892 m - >150 µm**

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.3
<i>Bulimina costata</i>	5	6								3.0
<i>Bulimina inflata</i>	24	16								11.0
<i>Bulimina marginata</i>	1									0.3
<i>Chilostomella ovoidea</i>							1			0.3
<i>Fissurina</i> sp.6	1									0.3
<i>Gavelinopsis translucens</i>	6	2								2.2
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>	1	3	1	1	1	2	3	1	5	5.0
<i>Gyroidina orbicularis</i>	1	1								0.6
<i>Oridorsalis tenera</i>	2	1								0.8
<i>Protolagena gracillima</i>			1							0.3
<i>Pullenia bulloides</i>	4	3	1							2.2
<i>Rutherfordoides mexicana</i>	4	1					1		1	1.9
<i>Uvigerina auberiana</i>	1	6	4							3.0
<i>Valvulineria laevigata</i>	2	12	1							4.1
<b>Porcelaneous taxa</b>										
<i>Pyrgo depressa</i>		3								0.8
<i>Pyrgo murrhina</i>		1								0.3
<i>Pyrgo serrata</i>	3									0.8
<i>Quinqueloculina seminula</i>		2								0.6
<i>Spiroloculina rotunda</i>	3	1								1.1
<i>Triloculina</i> spp.		1								0.3
<b>Agglutinated taxa</b>										
Agglutinated indet.		3								0.8
<i>Ammoscalaria</i> sp.	2									0.6
<i>Cyclammia</i> sp.	3	1								1.1
<i>Hormosina distans</i>	1			1						0.6
<i>Hormosina globulifera</i>		3								0.8
<i>Hormosina guttifera</i> type A	1									0.3
<i>Lagenammia</i> sp.1			3	1						1.1
<i>Lagenammia</i> sp.2							4			1.1
<i>Nouria harrisii</i>		7					12	2		5.8
<i>Psammosphaera</i> spp.		9								2.5
<i>Recurvoides contortus</i>	1	14		2						4.7
<i>Reophax bilocularis</i> type A	45	26								19.6
<i>Reophax pilulifer</i>	49	21	3		1					20.4
<i>Tetragonostomina rhombiformis</i>	2	3								1.4
<b>Standing Stocks (not normalized)</b>	162	147	14	5	2	2	21	3	6	100
<b>Simple Diversity</b>	22	25	7	4	2	1	5	2	2	

**Station 6 - 892 m - >150 µm**

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.3
<i>Bulimina costata</i>	5	6								3.0
<i>Bulimina inflata</i>	24	16								11.0
<i>Bulimina marginata</i>	1									0.3
<i>Chilostomella ovoidea</i>							1			0.3
<i>Fissurina</i> sp.6	1									0.3
<i>Gavelinopsis translucens</i>	6	2								2.2
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>	1	3	1	1	1	2	3	1	5	5.0
<i>Gyroidina orbicularis</i>	1	1								0.6
<i>Oridorsalis tenera</i>	2	1								0.8
<i>Protolagena gracillima</i>			1							0.3
<i>Pullenia bulloides</i>	4	3	1							2.2
<i>Rutherfordoides mexicana</i>	4	1					1		1	1.9
<i>Uvigerina auberiana</i>	1	6	4							3.0
<i>Valvulineria laevigata</i>	2	12	1							4.1
<b>Porcelaneous taxa</b>										
<i>Pyrgo depressa</i>		3								0.8
<i>Pyrgo murrhina</i>		1								0.3
<i>Pyrgo serrata</i>	3									0.8
<i>Quinqueloculina seminula</i>		2								0.6
<i>Spiroloculina rotunda</i>	3	1								1.1
<i>Triloculina</i> spp.		1								0.3
<b>Agglutinated taxa</b>										
Agglutinated indet.		3								0.8
<i>Ammoscalaria</i> sp.	2									0.6
<i>Cyclammia</i> sp.	3	1								1.1
<i>Hormosina distans</i>	1			1						0.6
<i>Hormosina globulifera</i>		3								0.8
<i>Hormosina guttifera</i> type A	1									0.3
<i>Lagenammia</i> sp.1			3	1						1.1
<i>Lagenammia</i> sp.2							4			1.1
<i>Nouria harrisii</i>		7					12	2		5.8
<i>Psammosphaera</i> spp.		9								2.5
<i>Recurvoides contortus</i>	1	14		2						4.7
<i>Reophax bilocularis</i> type A	45	26								19.6
<i>Reophax pilulifer</i>	49	21	3		1					20.4
<i>Tetragonostomina rhombiformis</i>	2	3								1.4
<b>Standing Stocks (not normalized)</b>	162	147	14	5	2	2	21	3	6	100
<b>Simple Diversity</b>	22	25	7	4	2	1	5	2	2	

**Station 5 - 1150 m - >150 µm**

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.1
<i>Bulimina costata</i>		1								0.1
<i>Bulimina elegans</i> var. <i>exilis</i>		1								0.1
<i>Bulimina inflata</i>	11	13								3.1
<i>Bulimina marginata</i>		1								0.1
<i>Cibicides</i> sp.1	12	1								1.7
<i>Cibicides refulgens</i>	17	34	9							7.7
<i>Cibicides wuellerstorfi</i>	1									0.1
<i>Cibicoides</i> sp.	1									0.1
<i>Cibicoides</i> sp.2		4								0.5
<i>Cibicoides kullenbergi</i>	124	45	5	1	1					22.6
<i>Dentalina</i> sp.									1	0.1
<i>Fissurina marginata</i>			1							0.1
<i>Gavelinopsis translucens</i>	13	7								2.6
<i>Globobulimina pyrula</i>	1				1					0.3
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>		1								0.1
<i>Globocassidulina subglobosa</i>		1								0.1
<i>Gyroidina neosoldanii</i>	1	4								0.6
<i>Gyroidina orbicularis</i>		4								0.5
<i>Hoeglundina elegans</i>			1							0.1
<i>Nonion</i> sp.						1				0.1
<i>Oridorsalis tenera</i>	1	2								0.4
<i>Protolagena gracillima</i>		1								0.1
<i>Pullenia bulloides</i>		5								0.6
<i>Rutherfordoides mexicana</i>		1			2	1	5	2		1.4
<i>Sphaeroidina bulloides</i>	6	8								1.8
<i>Uvigerina auberiana</i>		11								1.4
<i>Uvigerina peregrina</i>	1	1	1							0.4
<i>Valvulineria laevigata</i>	1	10	2							1.7
<b>Porcelaneous taxa</b>										
Miliolid indet.	2	1								0.4
<i>Biloculinella irregularis</i>	1	3								0.5
<i>Biloculinella labiata</i>		7								0.9
<i>Cornuspira</i> sp.		1								0.1
<i>Cornuspira involvens</i>	1									0.1
<i>Discospirina</i> sp.	2									0.3
<i>Pyrgo</i> sp.2	4									0.5
<i>Pyrgo depressa</i>	4									0.5
<i>Pyrgo elongata</i>	2	2								0.5
<i>Pyrgo lucemula</i>		1								0.1
<i>Pyrgo murrhina</i>	1									0.1
<i>Pyrgo serrata</i>	1									0.1
<i>Pyrgo verpestilio</i>	1									0.1
<i>Pyrgoella sphaera</i>		6								0.8
<i>Quinqueloculina</i> sp.	1									0.1
<i>Quinqueloculina badenensis</i>	1									0.1
<i>Quinqueloculina seminula</i>	11	3								1.8
<i>Triloculina</i> spp.	1	1								0.3
<b>Agglutinated taxa</b>										
Epilithic indet.		1								0.1
<i>Ammodiscus catinus</i>	1	1								0.3
<i>Ammolagena clavata</i>	1									0.1
<i>Ammoscalaria</i> sp.		2								0.3
<i>Cribrostomoides subglobosus</i>	8	8								2.1
<i>Cyclammina</i> sp.		3								0.4
<i>Eggerella bradyi</i>	42	31	3							9.8
<i>Haplophragmoides</i> sp.		1								0.1
<i>Hormosina globulifera</i>	3	12								1.9
<i>Lagenammina pseudodifflugiformis</i>		12	5	6	2	1				3.3
<i>Nouria harrisii</i>	1		3		4	3	13	7	10	5.3
<i>Recurvoides contortus</i>	29	29	3							7.8
<i>Reophax bilocularis</i> type A	19	47	4	2						9.2
<i>Reophax micaceus</i>							3			0.4
<i>Reophax nodulosus</i>	9									1.2
<i>Reophax pilulifer</i>		9					1			1.3
<i>Trochammina conica</i>		1								0.1
<b>Standing Stocks (not normalized)</b>	336	339	37	9	10	6	22	9	11	100
<b>Simple Diversity</b>	36	44	11	3	5	4	4	2	2	

**Station 4 - 1957 m - >150 µm**

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.1
Epilithic indet.		1								0.1
<i>Anomalinoidea</i> sp.		3								0.2
<i>Bolivina subspinescens</i>		1								0.1
<i>Bulimina elegans</i> var. <i>exilis</i>						1				0.1
<i>Bulimina inflata</i>	49	6	1							4.4
<i>Cassidulina laevigata</i> var. <i>carinata</i>	11	5								1.3
<i>Chilostomella ovoidea</i>	2	1					2		2	0.6
<i>Cibicides</i> sp.1	18	2								1.6
<i>Cibicides wuellerstorfi</i>	18	3								1.7
<i>Cibicides</i> sp.1	1	1								0.2
<i>Cibicides</i> sp.2	9	3	1							1.0
<i>Cibicides bradyi</i>	3	2								0.4
<i>Cibicides</i> ? <i>globulosus</i>	5									0.4
<i>Cibicides</i> cf. <i>pachydermus</i>	2									0.2
<i>Cibicides kullenbergi</i>	138	10	6	2			3	2	1	12.8
<i>Epistominella exigua</i>	6	5								0.9
<i>Fissurina</i> sp.5	1									0.1
<i>Gavelinopsis translucens</i>	38	5	1							3.5
<i>Globobulimina pyrula</i>						1	2	1		0.3
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>		3					1		1	0.4
<i>Globocassidulina subglobosa</i>	1									0.1
<i>Gyroldina neosoldanii</i>	35	41	12	6	2					7.6
<i>Gyroldina orbicularis</i>	7	10	1	1	2	1				1.7
<i>Lagenosolenia</i> sp.	1									0.1
<i>Lenticulina gibba</i>		1								0.1
<i>Melonis barleeanus</i>	8	8	9	13	21	7	11	4	4	6.7
<i>Nonion</i> sp.	1									0.1
<i>Nonionella atlantica</i>	1									0.1
<i>Oridorsalis tenera</i>	7	8								1.2
<i>Pullenia</i> sp.					2					0.2
<i>Pullenia bulloides</i>	10		1							0.9
<i>Pullenia quinqueloba</i>	2	1								0.2
<i>Robertina translucens</i>	1	1								0.2
<i>Rutherfordoidea mexicana</i>								1		0.1
<i>Sphaeroidina bulloides</i>	4	1								0.4
<i>Uvigerina auferiana</i>		1		1						0.2
<i>Uvigerina peregrina</i>	51	27	5				2			6.7
<i>Valvulineria laevigata</i>		3	3							0.5
<b>Porcelaneous taxa</b>										
<i>Biloculina irregularis</i>	4	1								0.4
<i>Pyrgo murrhina</i>	4									0.3
<i>Pyrgoella sphaera</i>	1									0.1
<i>Triloculina tricarinata</i>	1									0.1
<b>Agglutinated taxa</b>										
Indet.		3								0.2
Epilithic indet.	1									0.1
<i>Ammoglobigerina</i> sp.	4									0.3
? <i>Chitinosaccus</i> sp.	20	13					1			2.7
<i>Cribrostomoides subglobosus</i>		1								0.1
<i>Eggerella bradyi</i>	18	2					1			1.7
<i>Hormosina globulifera</i>		3								0.2
<i>Hormosina guttifer</i> type A		1								0.1
<i>Hormosina guttifer</i> type B	8									0.6
<i>Karriella chilostoma</i>	8	1	1							0.8
<i>Lagenammina difflugiiformis</i> var. <i>calcare</i> type B		1								0.1
<i>Lagenammina pseudodifflugiiformis</i>	3	1								0.3
<i>Psammosphaera</i> spp.	23	4								2.1
<i>Recurvoides contortus</i>	24	8								2.5
<i>Reophax bilocularis</i> type A	99	28								10.0
<i>Reophax bilocularis</i> type B	1									0.1
<i>Reophax dentaliniformis</i>	5	4								0.7
<i>Reophax fusiformis</i>		2								0.2
<i>Reophax micaceus</i>	5	1								0.5
<i>Reophax nodulosus</i>		3								0.2
<i>Reophax pilulifer</i>	10	5								1.2
<i>Reophax scoriurus</i>	4	5								0.7
<i>Tetragonostomina rhombiformis</i>	3									0.2
<i>Tholosina</i> sp.		1								0.1
<i>Trochammina</i> sp.	1									0.1
<i>Trochammina globigeriniformis</i>	73	17								7.1
<i>Trochammina nitida</i>	3									0.2
<i>Vanhoeffenella gaussi</i>	3									0.2
<b>Soft-shelled</b>										
Soft-shelled spp.	58	51								8.6
<b>Standing Stocks (not normalised)</b>	814	310	41	23	27	10	23	8	8	100
<b>Simple Diversity</b>	52	49	11	5	4	4	8	4	4	

**Station 3 - 2974 m - >150 µm**

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Indet.	2									0.2
<i>Bulimina inflata</i>	2									0.2
<i>Chilostomella ovoidea</i>	1									0.1
<i>Cibicides</i> sp.1	1									0.1
<i>Cibicides wuellerstorfi</i>	16	3								1.7
<i>Cibicidoides</i> sp.1	2	2								0.4
<i>Cibicidoides bradyi</i>	1	1								0.2
<i>Cibicidoides kullenbergi</i>	2	2								0.2
<i>Epistominella exigua</i>	2	5								0.6
<i>Fissurina</i> sp.1	1									0.1
<i>Fissurina</i> sp.2		1								0.1
<i>Fissurina</i> sp.3		1	1							0.2
<i>Fissurina</i> sp.4		1								0.1
<i>Fissurina carinata</i>	1	1								0.2
<i>Fissurina longispina</i>	1									0.1
<i>Fissurina marginata</i>	1									0.1
<i>Fissurina orbignyana</i>	1				1					0.2
<i>Globocassidulina subglobosa</i>	3									0.3
<i>Gyroidina neosoldanii</i>		4								0.4
<i>Gyroidina orbicularis</i>		2					1			0.3
<i>Gyroidina umbonata</i>	2									0.2
<i>Hoeglundina elegans</i>	10	9	5	1						2.2
<i>Melonis barleeanus</i>	1	4	1							0.5
<i>Oridorsalis tenera</i>	6	3								0.8
<i>Proterolagena gracillima</i>	2	2								0.4
<i>Pullenia quinqueloba</i>		3						1		0.4
<i>Uvigerina auberiana</i>		1								0.1
<i>Uvigerina peregrina</i>	6	7								1.2
<b>Porcelaneous taxa</b>										
Indet.		2								0.2
<i>Bilocalina irregularis</i>	11	4								1.3
<i>Cornuloculina inconstans</i>	1									0.1
<i>Miliolinella</i> sp.	1									0.1
<i>Ophthalmidium acutumargo</i>	2	1								0.3
<i>Pyrgo</i> sp.	1									0.1
<i>Pyrgo murrhina</i>	4	1			1					0.5
<i>Quinqueloculina badenensis</i>		1								0.1
<i>Quinqueloculina elongata</i>		1								0.1
<i>Quinqueloculina seminula</i>	1	1								0.2
<i>Sigmoilina tenuis</i>		1								0.1
<i>Spirillina</i> sp.					1					0.1
<i>Spiroloculina</i> sp.	1									0.1
<i>Triloculina trigonula</i>		1								0.1
<b>Agglutinated taxa</b>										
Indet.	1									0.1
<i>Adercotryma glomerata</i>	1	2								0.3
<i>Ammodiscus catinus</i>	7	2								0.8
<i>Ammolagena clavata</i>	1	1								0.2
<i>Ammoscalaria</i> sp.	1	2								0.3
? <i>Chitinosaccus</i> sp.								3		0.3
<i>Cribrostomoides subglobosus</i>	10						1			1.0
<i>Eggerella</i> sp.	1									0.1
<i>Eggerella bradyi</i>	4	4								0.7
<i>Haplophragmoides</i> sp.		1								0.1
<i>Hippocrepinella alba</i>	4	2								0.5
<i>Hormosina guttifer</i> type A	8	1		1						0.9
<i>Lagenammina difflugiformis</i> forma typica	29	14	5	3				1		4.6
<i>Lagenammina difflugiformis</i> var. calcarea type A	13	26								3.5
<i>Lagenammina difflugiformis</i> var. calcarea type B	27	16	1							3.9
<i>Lagenammina pseudodifflugiformis</i>	4	2								0.5
<i>Paratrochammina</i> ( <i>Paratrochammina</i> ) <i>earlandi</i>	16	2								1.6
<i>Recurvoides contortus</i>							1			0.1
<i>Reophax bilocularis</i> type A	519	160	2	1	1			1		60.5
<i>Reophax bilocularis</i> type B		3								0.3
<i>Reophax fusiformis</i>		3								0.3
<i>Reophax micaceus</i>	3	5								0.7
<i>Reophax nodulosus</i>	4	6								0.9
<i>Reophax pilulifer</i>	7	2								0.8
<i>Reophax scoriurus</i>	1									0.1
<i>Saccammina sphaerica</i>	2	1								0.3
<i>Sigmoilopsis schlumbergeri</i>	2	2								0.4
<i>Siphotextularia heterostoma</i>	1	2								0.3
<i>Subreophax aduncus</i>	1									0.1
<i>Textularia</i> sp.	1									0.1
<i>Thurammina</i> sp.1	2	3								0.4
<i>Thurammina albicans</i>	1									0.1
<i>Thurammina papillata</i>	2	2								0.4
<i>Vanhoeffenella gaussi</i>	1	1								0.2
<b>Soft-shelled</b>										
Soft-shelled type A	5									0.4
Soft-shelled spp.	1	2								0.3
<b>Standing Stocks (not normalized)</b>	764	332	15	6	4	0	3	6	0	100
<b>Simple Diversity</b>	59	53	6	4	4	0	3	4	0	

### Station 6 - 892 m - >150 µm

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.3
<i>Bulimina costata</i>	5	6								3.0
<i>Bulimina inflata</i>	24	16								11.0
<i>Bulimina marginata</i>	1									0.3
<i>Chilostomella ovoidea</i>							1			0.3
<i>Fissurina</i> sp.6	1									0.3
<i>Gavelinopsis translucens</i>	6	2								2.2
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>	1	3	1	1	1	2	3	1	5	5.0
<i>Gyroidina orbicularis</i>	1	1								0.6
<i>Oridorsalis tenera</i>	2	1								0.8
<i>Protolagena gracillima</i>			1							0.3
<i>Pullenia bulloides</i>	4	3	1							2.2
<i>Rutherfordoides mexicana</i>	4	1					1		1	1.9
<i>Uvigerina auberiana</i>	1	6	4							3.0
<i>Valvulineria laevigata</i>	2	12	1							4.1
<b>Porcelaneous taxa</b>										
<i>Pyrgo depressa</i>		3								0.8
<i>Pyrgo murrhina</i>		1								0.3
<i>Pyrgo serrata</i>	3									0.8
<i>Quinqueloculina seminula</i>		2								0.6
<i>Spiroloculina rotunda</i>	3	1								1.1
<i>Triloculina</i> spp.		1								0.3
<b>Agglutinated taxa</b>										
Agglutinated indet.		3								0.8
<i>Ammoscalaria</i> sp.	2									0.6
<i>Cyclammia</i> sp.	3	1								1.1
<i>Hormosina distans</i>	1			1						0.6
<i>Hormosina globulifera</i>		3								0.8
<i>Hormosina guttifera</i> type A	1									0.3
<i>Lagenammia</i> sp.1			3	1						1.1
<i>Lagenammia</i> sp.2							4			1.1
<i>Nouria harrisii</i>		7					12	2		5.8
<i>Psammosphaera</i> spp.		9								2.5
<i>Recurvoides contortus</i>	1	14		2						4.7
<i>Reophax bilocularis</i> type A	45	26								19.6
<i>Reophax pilulifer</i>	49	21	3		1					20.4
<i>Tetragonostomina rhombiformis</i>	2	3								1.4
<b>Standing Stocks (not normalized)</b>	162	147	14	5	2	2	21	3	6	100
<b>Simple Diversity</b>	22	25	7	4	2	1	5	2	2	

**Station 5 - 1150 m - >150 µm**

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.1
<i>Bulimina costata</i>		1								0.1
<i>Bulimina elegans</i> var. <i>exilis</i>		1								0.1
<i>Bulimina inflata</i>	11	13								3.1
<i>Bulimina marginata</i>		1								0.1
<i>Cibicides</i> sp.1	12	1								1.7
<i>Cibicides refulgens</i>	17	34	9							7.7
<i>Cibicides wuellerstorfi</i>	1									0.1
<i>Cibicoides</i> sp.	1									0.1
<i>Cibicoides</i> sp.2		4								0.5
<i>Cibicoides kullenbergi</i>	124	45	5	1	1					22.6
<i>Dentalina</i> sp.									1	0.1
<i>Fissurina marginata</i>			1							0.1
<i>Gavelinopsis translucens</i>	13	7								2.6
<i>Globobulimina pyrula</i>	1				1					0.3
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>		1								0.1
<i>Globocassidulina subglobosa</i>		1								0.1
<i>Gyroidina neosoldanii</i>	1	4								0.6
<i>Gyroidina orbicularis</i>		4								0.5
<i>Hoeglundina elegans</i>			1							0.1
<i>Nonion</i> sp.						1				0.1
<i>Oridorsalis tenera</i>	1	2								0.4
<i>Protolagena gracillima</i>		1								0.1
<i>Pullenia bulloides</i>		5								0.6
<i>Rutherfordoides mexicana</i>		1			2	1	5	2		1.4
<i>Sphaeroidina bulloides</i>	6	8								1.8
<i>Uvigerina auberiana</i>		11								1.4
<i>Uvigerina peregrina</i>	1	1	1							0.4
<i>Valvulineria laevigata</i>	1	10	2							1.7
<b>Porcelaneous taxa</b>										
Miliolid indet.	2	1								0.4
<i>Biloculinella irregularis</i>	1	3								0.5
<i>Biloculinella labiata</i>		7								0.9
<i>Cornuspira</i> sp.		1								0.1
<i>Cornuspira involvens</i>	1									0.1
<i>Discospirina</i> sp.	2									0.3
<i>Pyrgo</i> sp.2	4									0.5
<i>Pyrgo depressa</i>	4									0.5
<i>Pyrgo elongata</i>	2	2								0.5
<i>Pyrgo lucemula</i>		1								0.1
<i>Pyrgo murrhina</i>	1									0.1
<i>Pyrgo serrata</i>	1									0.1
<i>Pyrgo verpestilio</i>	1									0.1
<i>Pyrgoella sphaera</i>		6								0.8
<i>Quinqueloculina</i> sp.	1									0.1
<i>Quinqueloculina badenensis</i>	1									0.1
<i>Quinqueloculina seminula</i>	11	3								1.8
<i>Triloculina</i> spp.	1	1								0.3
<b>Agglutinated taxa</b>										
Epilithic indet.		1								0.1
<i>Ammodiscus catinus</i>	1	1								0.3
<i>Ammolagena clavata</i>	1									0.1
<i>Ammoscalaria</i> sp.		2								0.3
<i>Cribrostomoides subglobosus</i>	8	8								2.1
<i>Cyclammina</i> sp.		3								0.4
<i>Eggerella bradyi</i>	42	31	3							9.8
<i>Haplophragmoides</i> sp.		1								0.1
<i>Hormosina globulifera</i>	3	12								1.9
<i>Lagenammina pseudodifflugiformis</i>		12	5	6	2	1				3.3
<i>Nouria harrisii</i>	1		3		4	3	13	7	10	5.3
<i>Recurvoides contortus</i>	29	29	3							7.8
<i>Reophax bilocularis</i> type A	19	47	4	2						9.2
<i>Reophax micaceus</i>							3			0.4
<i>Reophax nodulosus</i>	9									1.2
<i>Reophax pilulifer</i>		9					1			1.3
<i>Trochammina conica</i>		1								0.1
<b>Standing Stocks (not normalized)</b>	336	339	37	9	10	6	22	9	11	100
<b>Simple Diversity</b>	36	44	11	3	5	4	4	2	2	

**Station 4 - 1957 m - >150 µm**

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.1
Epilithic indet.		1								0.1
<i>Anomalinoidea</i> sp.		3								0.2
<i>Bolivina subspinescens</i>		1								0.1
<i>Bulimina elegans</i> var. <i>exilis</i>						1				0.1
<i>Bulimina inflata</i>	49	6	1							4.4
<i>Cassidulina laevigata</i> var. <i>carinata</i>	11	5								1.3
<i>Chilostomella ovoidea</i>	2	1					2		2	0.6
<i>Cibicides</i> sp.1	18	2								1.6
<i>Cibicides wuellerstorfi</i>	18	3								1.7
<i>Cibicides</i> sp.1	1	1								0.2
<i>Cibicides</i> sp.2	9	3	1							1.0
<i>Cibicides bradyi</i>	3	2								0.4
<i>Cibicides</i> ? <i>globulosus</i>	5									0.4
<i>Cibicides</i> cf. <i>pachydermus</i>	2									0.2
<i>Cibicides kullenbergi</i>	138	10	6	2			3	2	1	12.8
<i>Epistominella exigua</i>	6	5								0.9
<i>Fissurina</i> sp.5	1									0.1
<i>Gavelinopsis translucens</i>	38	5	1							3.5
<i>Globobulimina pyrula</i>						1	2	1		0.3
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>		3					1		1	0.4
<i>Globocassidulina subglobosa</i>	1									0.1
<i>Gyroldina neosoldanii</i>	35	41	12	6	2					7.6
<i>Gyroldina orbicularis</i>	7	10	1	1	2	1				1.7
<i>Lagenosolenia</i> sp.	1									0.1
<i>Lenticulina gibba</i>		1								0.1
<i>Melonis barleeanus</i>	8	8	9	13	21	7	11	4	4	6.7
<i>Nonion</i> sp.	1									0.1
<i>Nonionella atlantica</i>	1									0.1
<i>Oridorsalis tenera</i>	7	8								1.2
<i>Pullenia</i> sp.					2					0.2
<i>Pullenia bulloides</i>	10		1							0.9
<i>Pullenia quinqueloba</i>	2	1								0.2
<i>Robertina translucens</i>	1	1								0.2
<i>Rutherfordoidea mexicana</i>								1		0.1
<i>Sphaeroidina bulloides</i>	4	1								0.4
<i>Uvigerina auferiana</i>		1		1						0.2
<i>Uvigerina peregrina</i>	51	27	5				2			6.7
<i>Valvulineria laevigata</i>		3	3							0.5
<b>Porcelaneous taxa</b>										
<i>Biloculina irregularis</i>	4	1								0.4
<i>Pyrgo murrhina</i>	4									0.3
<i>Pyrgoella sphaera</i>	1									0.1
<i>Triloculina tricarinata</i>	1									0.1
<b>Agglutinated taxa</b>										
Indet.		3								0.2
Epilithic indet.	1									0.1
<i>Ammoglobigerina</i> sp.	4									0.3
? <i>Chitinosaccus</i> sp.	20	13					1			2.7
<i>Cribrostomoides subglobosus</i>		1								0.1
<i>Eggerella bradyi</i>	18	2					1			1.7
<i>Hormosina globulifera</i>		3								0.2
<i>Hormosina guttifer</i> type A		1								0.1
<i>Hormosina guttifer</i> type B	8									0.6
<i>Karriella chilostoma</i>	8	1	1							0.8
<i>Lagenammina difflugiformis</i> var. <i>calcare</i> type B		1								0.1
<i>Lagenammina pseudodifflugiformis</i>	3	1								0.3
<i>Psammosphaera</i> spp.	23	4								2.1
<i>Recurvoides contortus</i>	24	8								2.5
<i>Reophax bilocularis</i> type A	99	28								10.0
<i>Reophax bilocularis</i> type B	1									0.1
<i>Reophax dentaliniformis</i>	5	4								0.7
<i>Reophax fusiformis</i>		2								0.2
<i>Reophax micaceus</i>	5	1								0.5
<i>Reophax nodulosus</i>		3								0.2
<i>Reophax pilulifer</i>	10	5								1.2
<i>Reophax scoriurus</i>	4	5								0.7
<i>Tetragonostomina rhombiformis</i>	3									0.2
<i>Tholosina</i> sp.		1								0.1
<i>Trochammina</i> sp.	1									0.1
<i>Trochammina globigeriniformis</i>	73	17								7.1
<i>Trochammina nitida</i>	3									0.2
<i>Vanhoeffenella gaussi</i>	3									0.2
<b>Soft-shelled</b>										
Soft-shelled spp.	58	51								8.6
<b>Standing Stocks (not normalised)</b>	814	310	41	23	27	10	23	8	8	100
<b>Simple Diversity</b>	52	49	11	5	4	4	8	4	4	



### Station 6 - 892 m - >150 µm

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.3
<i>Bulimina costata</i>	5	6								3.0
<i>Bulimina inflata</i>	24	16								11.0
<i>Bulimina marginata</i>	1									0.3
<i>Chilostomella ovoidea</i>							1			0.3
<i>Fissurina</i> sp.6	1									0.3
<i>Gavelinopsis translucens</i>	6	2								2.2
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>	1	3	1	1	1	2	3	1	5	5.0
<i>Gyroidina orbicularis</i>	1	1								0.6
<i>Oridorsalis tenera</i>	2	1								0.8
<i>Protolagena gracillima</i>			1							0.3
<i>Pullenia bulloides</i>	4	3	1							2.2
<i>Rutherfordoides mexicana</i>	4	1					1		1	1.9
<i>Uvigerina auberiana</i>	1	6	4							3.0
<i>Valvulineria laevigata</i>	2	12	1							4.1
<b>Porcelaneous taxa</b>										
<i>Pyrgo depressa</i>		3								0.8
<i>Pyrgo murrhina</i>		1								0.3
<i>Pyrgo serrata</i>	3									0.8
<i>Quinqueloculina seminula</i>		2								0.6
<i>Spiroloculina rotunda</i>	3	1								1.1
<i>Triloculina</i> spp.		1								0.3
<b>Agglutinated taxa</b>										
Agglutinated indet.		3								0.8
<i>Ammoscalaria</i> sp.	2									0.6
<i>Cyclammia</i> sp.	3	1								1.1
<i>Hormosina distans</i>	1			1						0.6
<i>Hormosina globulifera</i>		3								0.8
<i>Hormosina guttifera</i> type A	1									0.3
<i>Lagenammia</i> sp.1			3	1						1.1
<i>Lagenammia</i> sp.2							4			1.1
<i>Nouria harrisii</i>		7					12	2		5.8
<i>Psammosphaera</i> spp.		9								2.5
<i>Recurvoides contortus</i>	1	14		2						4.7
<i>Reophax bilocularis</i> type A	45	26								19.6
<i>Reophax pilulifer</i>	49	21	3		1					20.4
<i>Tetragonostomina rhombiformis</i>	2	3								1.4
<b>Standing Stocks (not normalized)</b>	162	147	14	5	2	2	21	3	6	100
<b>Simple Diversity</b>	22	25	7	4	2	1	5	2	2	

**Station 5 - 1150 m - >150 µm**

Depth interval in the sediment (cm)	0-0.5	0.5-1.0	1.0-1.5	1.5-2.0	2.0-2.5	2.5-3.0	3.0-3.5	3.5-4.0	4.0-5.0	%
<b>Foraminiferal taxa</b>										
<b>Perforate calcareous taxa</b>										
Perforate indet.		1								0.1
<i>Bulimina costata</i>		1								0.1
<i>Bulimina elegans</i> var. <i>exilis</i>		1								0.1
<i>Bulimina inflata</i>	11	13								3.1
<i>Bulimina marginata</i>		1								0.1
<i>Cibicides</i> sp.1	12	1								1.7
<i>Cibicides refulgens</i>	17	34	9							7.7
<i>Cibicides wuellerstorfi</i>	1									0.1
<i>Cibicoides</i> sp.	1									0.1
<i>Cibicoides</i> sp.2		4								0.5
<i>Cibicoides kullenbergi</i>	124	45	5	1	1					22.6
<i>Dentalina</i> sp.									1	0.1
<i>Fissurina marginata</i>			1							0.1
<i>Gavelinopsis translucens</i>	13	7								2.6
<i>Globobulimina pyrula</i>	1				1					0.3
<i>Globobulimina pyrula</i> var. <i>pseudospinescens</i>		1								0.1
<i>Globocassidulina subglobosa</i>		1								0.1
<i>Gyroidina neosoldanii</i>	1	4								0.6
<i>Gyroidina orbicularis</i>		4								0.5
<i>Hoeglundina elegans</i>			1							0.1
<i>Nonion</i> sp.							1			0.1
<i>Oridorsalis tenera</i>	1	2								0.4
<i>Protolagena gracillima</i>		1								0.1
<i>Pullenia bulloides</i>		5								0.6
<i>Rutherfordoides mexicana</i>		1			2	1	5	2		1.4
<i>Sphaeroidina bulloides</i>	6	8								1.8
<i>Uvigerina auberiana</i>		11								1.4
<i>Uvigerina peregrina</i>	1	1	1							0.4
<i>Valvulineria laevigata</i>	1	10	2							1.7
<b>Porcelaneous taxa</b>										
Miliolid indet.	2	1								0.4
<i>Biloculinella irregularis</i>	1	3								0.5
<i>Biloculinella labiata</i>		7								0.9
<i>Cornuspira</i> sp.		1								0.1
<i>Cornuspira involvens</i>	1									0.1
<i>Discospirina</i> sp.	2									0.3
<i>Pyrgo</i> sp.2	4									0.5
<i>Pyrgo depressa</i>	4									0.5
<i>Pyrgo elongata</i>	2	2								0.5
<i>Pyrgo lucemula</i>		1								0.1
<i>Pyrgo murrhina</i>	1									0.1
<i>Pyrgo serrata</i>	1									0.1
<i>Pyrgo verpestilio</i>	1									0.1
<i>Pyrgoella sphaera</i>		6								0.8
<i>Quinqueloculina</i> sp.	1									0.1
<i>Quinqueloculina badenensis</i>	1									0.1
<i>Quinqueloculina seminula</i>	11	3								1.8
<i>Triloculina</i> spp.	1	1								0.3
<b>Agglutinated taxa</b>										
Epilithic indet.		1								0.1
<i>Ammodiscus catinus</i>	1	1								0.3
<i>Ammolagena clavata</i>	1									0.1
<i>Ammoscalaria</i> sp.		2								0.3
<i>Cribrostomoides subglobosus</i>	8	8								2.1
<i>Cyclammina</i> sp.		3								0.4
<i>Eggerella bradyi</i>	42	31	3							9.8
<i>Haplophragmoides</i> sp.		1								0.1
<i>Hormosina globulifera</i>	3	12								1.9
<i>Lagenammina pseudodifflugiformis</i>		12	5	6	2	1				3.3
<i>Nouria harrisii</i>	1		3		4	3	13	7	10	5.3
<i>Recurvoides contortus</i>	29	29	3							7.8
<i>Reophax bilocularis</i> type A	19	47	4	2						9.2
<i>Reophax micaceus</i>							3			0.4
<i>Reophax nodulosus</i>	9									1.2
<i>Reophax pilulifer</i>		9					1			1.3
<i>Trochammina conica</i>		1								0.1
<b>Standing Stocks (not normalized)</b>	336	339	37	9	10	6	22	9	11	100
<b>Simple Diversity</b>	36	44	11	3	5	4	4	2	2	